



Review

Review on the use of AI-based methods and tools for treating mental conditions and mental rehabilitation

Vladimir Khorev^{1,2,3,a} , Anton Kiselev^{1,b}, Artem Badarin^{1,2,c}, Vladimir Antipov^{1,2,d}, Oxana Drapkina^{1,e}, Semen Kurkin^{2,f}, and Alexander Hramov^{2,g}

¹ Coordinating Center for Fundamental Research, National Medical Research Center for Therapy and Preventive Medicine, 10 Petroverigsky per., Moscow 101990, Russia

² Baltic Center for Neurotechnology and Artificial Intelligence, Immanuel Kant Baltic Federal University, 14 Alexander Nevsky Street, Kaliningrad 236016, Russia

³ Neuroscience and Cognitive Technology Laboratory, Innopolis University, 1 Universitetskaya Street, Innopolis 420500, Russia

Received 25 June 2024 / Accepted 1 August 2024 / Published online 12 August 2024

© The Author(s), under exclusive licence to EDP Sciences, Springer-Verlag GmbH Germany, part of Springer Nature 2024

Abstract This review provides a thorough examination of recent developments in artificial intelligence analysis methods within mental and psychiatry field. By analyzing and comparing results obtained with various tools and techniques, we provide a comprehensive and systematic understanding of applications. Our main methods include meta-analysis, search queries with the keywords and network-based approach. In our analysis, we observed that terms associated with robotics, human–computer interaction, speech perception, and certain applications, such as chronic fatigue syndrome and psychological adaptation, have been gradually losing prominence. And conversely, techniques such as deep learning, virtual reality, and virtual assistance are gaining traction, and increasing interest was noted for applications involving autistic spectrum disorders, mild cognitive impairments, and psychiatric research areas. The structured and organized presentation of information, along with the accompanying visualizations and diagrams, makes it a valuable resource for scientists and researchers working in the domains of artificial intelligence.

Abbreviations

Add/adhd	Attention-deficit disorder or attention-deficit hyperactivity disorder
AI	Artificial intelligence
AR	Augmented reality
ASD	Autism spectrum disorder
BCI	Brain–computer interface
CBT	Cognitive and behavioral therapy
CNN	Convolutional neural network
CVA	Computer virtual assistant or chatbot
DL	Deep learning
FC	Functional connectivity
MCI	Mild cognitive impairments
II	Image interpretation
ML	Machine learning

^a e-mail: khorevvs@gmail.com (corresponding author)

^b e-mail: kiselev@gnicpm.ru

^c e-mail: badarin.a.a@mail.ru

^d e-mail: vantipovm@gmail.com

^e e-mail: drapkina@bk.ru

^f e-mail: kurkinsa@gmail.com

^g e-mail: hramovae@gmail.com

NLP	Natural language processing
OCD	Obsessive-compulsive disorder
PTSD	Post-traumatic stress disorder
SVM	Support vector machine
VG	Video games
VR	Virtual reality
XAI	Explainable artificial intelligence

1 Introduction

Artificial intelligence (AI) in psychiatry has become increasingly prominent as a tool for diagnosis [1–5], treatment planning [6–9], and patient monitoring [10–13]. AI technologies, such as machine learning algorithms and natural language processing, are being integrated into psychiatric practice to enhance the accuracy and efficiency of psychiatric assessments, predict disease progression, and personalize treatment plans. In addition, AI-driven virtual assistants are being explored for their potential in providing remote mental health support.

Among mental health patients, AI can be particularly helpful for:

- Individuals with complex mental health disorders, where AI-driven tools can improve the accuracy of diagnosis and enhance personalized treatment plans for conditions such as schizophrenia [14–18], bipolar disorder [19–21], severe depression [22–28], autism spectrum disorder [29, 30];
- Treatment-resistant conditions: AI algorithms can help identify patterns in patient data to suggest novel treatment approaches or combinations of medications for individuals who do not respond to traditional therapies [31–37];
- Remote or hard-to-reach populations: AI-powered virtual assistants and telehealth platforms can increase access to mental health services for individuals who live in remote areas or face mobility challenges [38–41].

These advances span across various fields, including:

- Psychiatry and psychology: improving diagnostic accuracy and personalized treatment planning for mental health disorders [42, 43];
- Neuroscience and cognitive psychology: enhancing understanding of brain functioning and cognitive processes [44–46];
- Computational and data sciences: developing advanced AI algorithms for data analysis, pattern recognition, and predictive modeling [47–51].

There is an increasing use of AI for addressing issues in the fields of psychology and psychiatry, suggesting a trend towards more widespread adoption of this technology in mental health care (see Fig. 1). By examining the uses of AI in the treatment of various mental disorders, it is possible provide insight into the potential benefits and limitations of this technology and its usefulness for improving clinical outcomes. AI-based methods can be used for more precise adjustment of psychological testing, taking into account the individual mental characteristics of patients [52–54].

A large number of systematic reviews on AI in medicine have been created, which tend to have a specific and limited focus on solving a particular issue or addressing a specific problem [40, 55–64]. Existing works cover a broader range of AI applications and their effectiveness across different patient populations. But meta-reviews on AI in mental rehabilitation are important because they provide a comprehensive assessment of the current state of research and application in this domain. Such reviews can help identify trends, gaps, and future directions in the research by synthesizing findings across multiple studies. They can also highlight methodological limitations and ethical concerns in existing studies and thus guide future research and practice in the field. In addition, meta-reviews can provide an objective evaluation of the efficacy and effectiveness of AI-based interventions in mental rehabilitation, which is crucial for informing evidence-based decision-making.

Certain gaps where AI integration in psychiatry warrants further review and exploration include:

- Limited ethical guidelines for AI-driven mental health services, particularly regarding data privacy and consent [65–71];
- Limited generalizability of AI-powered tools across different populations and cultures;
- Need for user-friendly AI applications that integrate smoothly into clinical workflows without disrupting established care models;

- Further research into the specific benefits of AI-driven mental health services, in terms of cost-effectiveness and improved patient outcomes compared to traditional approaches.

In this review, we aim to explore the tendencies of using AI in psychiatry from data science perspective. We identify and compare technologies for different applications used by researchers in different branches of medicine. We analyzed the results obtained from PubMed database using thoroughly constructed search queries based on keywords to obtain the most conclusive results with each query.

2 Materials and methods

As part of the methodology implementation, published articles were retrieved from scientific databases for the last 30 years using PubMed search engine (<https://pubmed.ncbi.nlm.nih.gov>, accessed on 20 May 2024). The following terms were used in the construction of search queries to conduct an analytical review of the current scientific and technical, regulatory, methodological literature addressing the scientific and technical problem of AI for the rehabilitation, as well as the prevention of cognitive disorders and diagnosis of patients with motor or cognitive disorders within the framework of rehabilitation medicine: (“AI” or “Artificial Intelligence” or “Machine Learning”) and “rehabilitation” and (“cognitive” or “mental disorder” or “mental impairment” or “neurodegeneration” or “cognitive impairment”) keywords were mandatory part of the query, while terms for the methods (e.g., “Virtual reality”), and application (e.g., “dementia”) were combined pairwise as shown in Table 1.

At the initial stage of our review, we have identified and highlighted the following key issues that we wish to address:

- Which types of AI methods are the most frequently utilized for treating mental conditions?
- Are there varying preferences for them in different areas of medicine? If so, what are the reasons behind these discrepancies?
- which types of AI methods correspond to which fields of application?

During the research phase, we carefully reviewed papers that met the criteria of having one of the search queries or equivalent restatements in their title, abstract, or key words. In addition, we excluded papers with the types “Book”, “Chapter”, and “Monograph” from our final sample using PubMed search filters. Following the collection of the number of relevant papers found for each query, we chose the most significant results and features, and then performed a more specific review to identify the causes behind these findings on the basis of specific examples.


We created a keyword co-occurrence network which used information gathered from around 700 papers found in PubMed by using a search query using VOSviewer version 1.6.20 (Center for Science and Technology Studies, Leiden University, The Netherlands) [72]. This network provides a visual representation of the frequency with which certain terms appeared together in the studies analyzed, offering a clearer understanding of the overarching trends and connections that exist within the field. The methodology of VOSviewer visualization technique is provided in details in Ref. [73]. We have previously used a similar network-based approach to analyze AI methods in biomedical research [74].

By employing network visualization techniques, it became possible for us to detect clusters or communities within the keyword co-occurrence network by assigning a color code to each item based on the cluster to which it belongs. In our research, we utilized the VOSviewer algorithm which employs a modularity function to identify these groups [75]. The extracted clusters offer further information and insights regarding the relationship between keywords within the field being studied.

Table 1 List of the used keywords

AI methods and AI-based techniques	Application
Brain–computer interface; deep learning; functional connectivity; image interpretation; machine learning; robotics; virtual reality; natural language processing; support vector machine; computer/virtual assistant/chatbot; video games;	Aging; neurodegenerative diseases; add or adhd or attention-deficit disorder or attention-deficit hyperactivity disorder; ASD or autism spectrum disorders; stroke; depression; anxiety; Alzheimer; aphasia; bipolar disorder; dementia; mci or mild cognitive impairments; OCD or obsessive–compulsive disorder; Parkinson disease; psychotherapy; post-traumatic stress disorder or PTSD; schizophrenia

Fig. 1 Amount of relevant publications (including (“AI” or “Artificial Intelligence” or “Machine Learning”) and (“rehabilitation” and (“cognitive” or “mental disorder” or “mental impairment” or “neurodegeneration” or “cognitive impairment”) keywords) over years



3 Review

3.1 Fundamentals of AI

By artificial Intelligence, we understand methods and algorithms, systems, or models that enable machines to perform tasks that typically require human-like intelligence, such as learning [76], problem-solving [77], decision-making [68, 71, 78–80], and perception [81–83]. AI methods encompass a wide range of techniques, including machine learning, natural language processing, computer vision, robotics, and expert systems.

The basic principle behind developing AI is to create algorithms and programs that can mimic human intelligence and problem-solving capabilities. This is achieved through various techniques, including machine learning and deep learning methods, which enable AI systems to improve and adapt based on experience and feedback. Other principles include data-driven approaches, where algorithms are trained on large datasets to identify patterns and make predictions, as well as symbolic or knowledge-based approaches, which incorporate symbolic representations of domains and prior knowledge to enable more informed reasoning and decision-making.

An artificial intelligence system can contribute to new scientific understanding through three main avenues. First, by functioning as a “computational microscope”, it can furnish insights unattainable through conventional experimental methods. Second, by acting as a “resource of inspiration”, it can augment human imagination and innovation, thereby broadening the scope of scientific possibilities. The third dimension in which AI contributes to scientific understanding is its role as an “agent of understanding”. By substituting humans in the process of generalizing observations and transferring these novel scientific concepts to different phenomena, and crucially, conveying these insights to human scientists, AI enables a deeper and broader comprehension of complex scientific matters [84].

3.2 AI-based methods classification

Some common primary methods include machine learning (ML), deep learning (DL), natural language processing (NLP), computer vision, robotics, and expert systems. Machine learning involves training algorithms on large datasets, enabling them to learn patterns and make predictions or decisions based on new data [85, 86]. Deep learning is a subset of machine learning that employs artificial neural networks, inspired by biological neurons, to learn patterns and make predictions [87, 88]. NLP focuses on understanding, generating, and interacting with human language [89–91], while computer vision deals with understanding and processing visual information [92, 93]. Robotics focuses on the development of autonomous robots [94, 95], while expert systems employ rule-based logic programming to mimic the decision-making capability of human experts [96, 97]. In artificial intelligence and machine learning, the Support Vector Machine (SVM) is a supervised learning model that can be used for classification and regression analysis [98–102]. The SVM algorithm essentially constructs a hyperplane or a set of hyperplanes in a high-dimensional space to identify the class labels of new instances or to map the input features into an N -dimensional space.

Brain functional connectivity (FC) [103] for AI refers to the analysis of interactions and relationships between different regions of a brain or neural network to understand how they work together. Specifically, it involves measuring changes in the synchronization or phase relationship of signals between regions to identify patterns and correlations between regions and their associated functions [104–107]. This technique has been used for AI




Fig. 3 The co-occurrence network constructed using VOSviewer. The sizes of the nodes are determined by the weight of the corresponding item that indicate the importance of the item, and the color is determined by the publishing activity per year (blue color denotes nodes with the majority of publications of 2014 year, and yellow denotes nodes where corresponding publications are from 2023 to 2024)

Looking at the current trends in the field, presented in the Fig. 3, we can see that among the clusters under consideration, there is a relatively even distribution of more and less recently popular terms. Nevertheless, we can clearly see that terms related to robotics, user-computer interface, speech perception, and among applications—chronic fatigue syndrome and psychological adaptation—are gradually losing popularity. Conversely, techniques related to deep learning, VR, virtual assistance are gaining popularity, and trending applications include autistic spectrum disorders, mild cognitive impairments, and psychiatry.

Main tables for the various keyword combinations is presented in the Fig. 4. As a first interesting observation from our review, we can note the varying popularity of different combinations of methods and applications. Of all the combinations considered, the following should be noted as the most popular: stroke and robotics [9, 118–124] or CVA [7, 63, 125–127], dementia and FC [128–134] or SVM [102, 135, 136], aging and FC [97, 137–140]. About one-third of the possible combinations have no publications, and of the remaining, about half are represented by only one or two publications. Among the most obvious trends is the prevalence of assistive or robotic techniques for stroke rehabilitation. In general, the most popular application areas are stroke, aging, Alzheimer's, dementia, and MCI. The least represented are works related to PTSD, OCD, bipolar disorder. In terms of solution options, FC, SVM, CVA are the most pronounced, NLP and video games the least, apparently due to the complexity and cost of development. Looking at BCI and robotic solutions, it can be seen that they are most prevalent in cases where the diseases are characterized by motor impairments and do not affect diseases of a purely psychiatric nature. At the same time, DL and FC methods are common in relatively equal measure for most applications. Image interpretation has a similar scope to BCI, except for the additional extension to neurodegenerative diseases. VR is ubiquitous except for OCD, bipolar disorder and aphasia. NLP is not a widespread technique, but does include applications against aphasia and neurodegenerative disorders. SVM has applications everywhere except OCD, PTSD, psychiatry. CVA is the second most popular technique after FC, but also does not cover OCD, PTSD, instead it is presented for psychotherapy and schizophrenia. Video games one of the least common methods which, however, has a niche application against MCI and add/adhd.

Further, we highlight the most interesting developments for the found combinations.




Fig. 4 Amount of publications by application/data keywords. *BCI* brain–computer interface, *DL* deep learning, *FC* functional connectivity, *II* image interpretation, *ML* machine learning, *Rob* robotics, *VR* virtual reality, *NLP* natural language processing, *SVM* support vector machine, *CVA* computer/virtual assistant/chatbot, *VG* video games, *add/adhd* attention-deficit hyperactivity disorder, *ASD* autism spectrum disorder, *OCD* obsessive–compulsive disorder, *mci* mild cognitive impairments, *PTSD* post-traumatic stress disorder. The blue histograms provide an overview of the average $\pm$ SD number of publications related to each technique or application, respectively

3.3.1 AI interpretation techniques and functional connectivity

Better understanding means better prognosis and treatment. The process of identifying different subtypes of neurodegenerative diseases and analyzing the distinct stages of each subtype’s progression is crucial. By studying the underlying mechanisms and identifying specific biomarkers, researchers can better understand the course and trajectory of these diseases, leading to improved diagnostic methods and targeted therapies [141]. Functional connectivity of the brain at rest after the stroke emerges as a crucial predictor of responsiveness to treatment, demonstrating its importance, both individually and in combination with other patient-related aspects [142]. One innovative approach is Neuropsychological Assessment of Patients with Acquired Brain Injury: A Cluster Analysis Approach to Tackle Heterogeneity in Web-Based Cognitive Rehabilitation [143]. This study used cluster analysis to group individuals with acquired brain injury into different subgroups based on their neuropsychological profiles and utilized web-based cognitive rehabilitation tailored to the specific needs of each subgroup. This approach aimed to improve the efficacy of cognitive rehabilitation programs by addressing individual differences and improving treatment outcomes.

3.3.2 Natural language processing

The application of conversational agents equipped with unconstrained natural language input capabilities for health-related purposes is an emerging area of research. However, the few studies published in this field mostly employed quasi-experimental designs and rarely examined efficacy or safety. To advance the field, future studies would benefit from more robust experimental designs and standardized reporting [91, 144]. Using machine learning models, the authors one of the works compared the performance of features to detect dementia, overall and during time periods relative to dementia case ascertainment in health administrative database. They managed to identified dementia-related features by applying published lists, clinician input, and NLP with word embeddings to free-text progress and consult notes and organized features into thematic groups [145].

3.3.3 Machine learning and support vector machines

Research has revealed that while behavioral factors, multimodal neuroimaging data, and demographic information collectively contribute important information in predicting responses to rehabilitation for chronic aphasia following

a stroke [142]. One research [14] presented the implementation of a human-in-the-loop machine learning framework applied to an automated story recall assessment model. The evaluation of human verbal memory is a crucial aspect in determining neurocognitive function in psychiatry, and there are similarities to the anamnesis process, which involves gathering information about a patient history. A scoping review was conducted to summarize the range of existing evidence on how participation is captured and operationalized in pediatric re/habilitation research using AI-based assessment approaches [146]. A narrative review explored the applications of machine learning models in stroke rehabilitation, focusing on models such as random forest, logistic regression, and deep neural networks [63].

3.3.4 BCI and neurofeedback

One of the works proposed a game-based neurofeedback training system designed to improve cognitive performance in healthy elderly individuals and those suffering from amnesic mild cognitive impairment [147]. Specific focus should be given to combined approaches, such as bolstering the effects of non-invasive neuromodulation and closed-loop methodologies. By search of multimodal biomarkers and electric field modeling it is possible to guide targeting and quantify dosages of treatment with the power of machine learning algorithms to integrate data gathered, ultimately pinpointing clinical outcomes [148]. Interactive games with the biofeedback were successfully used for the rehabilitation tasks [123, 149, 150]. Studies have demonstrated that it is possible to enhance artificial agents' ability to anticipate human decisions in situations involving coordination issues through the integration of well-known solutions referred to as focal points into machine learning algorithms [151].

3.3.5 Chatbots and virtual assistants

A more innovative trend is the utilization of CVA powered by sophisticated artificial intelligence. A randomized controlled trial examined the effects of an entirely automated conversational agent that reinforced cognitive behavioral therapy (CBT) techniques through a text-message format, comparing it to an information-only control [25, 152–154]. The study found a significantly greater reduction in depressive symptoms among individuals in the intervention group [155]. Noteworthy uses of CVA include predicting lithium treatment effectiveness in individuals experiencing their first manic episode and customizing treatment plans for bipolar disorder by targeting neuroimaging-based treatment markers [156]. The quantitative analysis measured the CVA impact by comparing the average improvement in symptoms of depression between more and less active CVA service users. The qualitative analysis measured the app engagement and experience by analyzing in-app user feedback and evaluated the performance of a machine learning classifier to detect user objections during conversations [157]. Utilizing the mimicking of body movements by a virtual agent, it is possible to enhance synchronization behavior and rapport in individuals suffering from schizophrenia [158].

Preliminary research studies showcased a digital behavioral aid that gathers and reports quantitative data. This system comprises customizable smartglasses that deliver personalized coaching experiences via gamified augmented-reality applications powered by artificial intelligence for autism spectrum disorder. These applications offer coaching in areas such as emotion recognition, facing gaze, eye contact, and behavioral self-regulation, benefiting both children and adults [159, 160].

Loneliness is increasingly recognized as a significant public health issue that increases the risk of morbidity and mortality. Artificial agents, such as robots, conversational agents, and chatbots, represent an innovative approach in healthcare delivery and have been shown to alleviate patient loneliness by providing social support [161, 162]. Nevertheless, similar to doctor-patient relationships, the quality of the connection an individual has with an artificial agent can considerably influence its support efficacy and involvement in care provision [163].

3.3.6 Virtual reality

A review article [61] examines the application of virtual reality (VR) technology for cognitive rehabilitation, focusing on individuals with neurological conditions like stroke, traumatic brain injury, and neurodegenerative diseases. The review evaluates the effectiveness of VR-based interventions in improving cognitive functions, such as attention, memory, and problem-solving skills, in these populations. Acknowledging the presence of neuromuscular impairments, strategies such as kinetic guidance, combined with the use of virtual reality (VR) and augmented reality (AR) in interactive exercise programs, have been effectively implemented during gait training to enhance rehabilitation outcomes [164]. There is known experimental works involved a simple, target-directed reaching task, which focused on using sensory prediction errors rather than relying on cognitive strategies [165]. Virtual reality-based therapies hold significant potential to enhance both motor and functional skills in diverse age groups, particularly in cases of multiple sclerosis and other impairments, through promoting cortex reorganization and activating multiple neuronal connections [56, 166]. A narrative review was performed on the rehabilitation of acquired cognitive disorders, examining the current state of research and various approaches used in this field [61].

3.3.7 Deep learning

Deep learning is a subset of machine learning that uses artificial neural networks to model and analyze complex patterns and relationships in data. It is inspired by the structure and function of the brain, with multiple layers of interconnected nodes or neurons. Among the reported advances, there is a system which employed advanced machine learning and signal processing techniques to identify clinical sense from digital footprints, leveraging smartphone's native sensors in order to recognize suicide risk [167]. The systematic review on the broader scale [168] included 4 full-scale randomized controlled trials, along with feasibility, and quasi-experimental studies. The interventions employed various designs and targeted different mental health issues while utilizing a variety of therapeutic approaches. All included studies reported reductions in psychological distress postintervention. A positive effect was observed from using wearable digital AI intervention to improve socialization skills in children diagnosed with autism spectrum disorder [169]. Employing digital treatment as a therapeutic approach has shown promising outcomes in improving the executive functions, particularly inhibitory control and visuospatial working memory, in individuals with ADHD [170]. One study proposed a cognitive prosthesis device designed for addressing face memory impairment as a proof-of-concept for developing domain-specific cognitive prosthetic [171]. Among more simplified approaches: an artificial intelligence algorithm runs on a smartwatch and recognizes minutes of social interaction based on vocal features extracted from ambient audio samples, without relying on natural language processing [172]. A perspective review was conducted to explore neurotechnological solutions for addressing post-traumatic stress disorder (PTSD) [173]. Note also that DL may find application in reconstructing FC in various diseases of the nervous system and aging [174–176].

3.3.8 Robotics

A recent systematic review examined the efficacy and effectiveness of robotic rehabilitation in improving cognitive functioning [177]. Useful intellectual interaction system is founded on a foundation of cognitive psychology principles. With an understanding of educational robots and their features, it integrates traditional teaching activities synergistically. To achieve this, a robot-assisted talent training mode is designed, blending the strengths of educational robots with conventional teaching methodologies to optimize learning outcomes [178]. A robot-assisted visuomotor wrist training regiment leads to improvements in proprioceptive acuity and movement accuracy in the opposite wrist [179]. Robot-assisted somatosensory-based training involving progressively precise active wrist movements enhances both proprioceptive sensitivity and motor function in individuals experiencing strokes [118]. There are known reviews for current knowledge about social and service robots for elderly care [139, 180, 181]. Machine learning tools for identifying different types of motor activity, both real and imagined, of patients from video signals [182], myography [183], MEG- [184, 185], EEG- [186–188] and fNIRS-registration [189–191] of brain activity are of great importance here for diagnostics and rehabilitation [192–194].

3.3.9 Video games

In the realm of video games, a notable approach is the personalized Web-based cognitive rehabilitation for patients with ischemic stroke, where customized digital interventions are employed to enhance cognitive abilities and recovery in those affected by ischemic stroke or cognitive impairments [195–199]. An AI-based training game has been developed that can adjust the difficulty level automatically based on an individual's performance in order to improve their voluntary engagement and ultimately affect clinical outcomes [139, 200]. A scoping review on the role of AI in serious games and gamification for health is also published very recently [201].

4 Discussion

The field of AI in mental rehabilitation seems to have its own research traditions that define its strengths and weaknesses.

The modularity observed in the topic clusters Fig. 2 is likely due to the different focuses and subtopics within the broader field of mental rehabilitation using AI. The first cluster focuses on the technical aspects of AI, specifically machine learning methods, and their application in neurodegenerative diseases that require significant support. The second cluster highlights AI's role in cognitive impairments and rehabilitative methods, such as memory enhancement and attention training. The third cluster encompasses psychological and psychiatric factors, such as motivation, quality of life, anxiety, and depression, which can significantly impact mental rehabilitation.

These factors can significantly impact the effectiveness of mental health and rehabilitation interventions. Interestingly, one study [202] mentioned an interesting and valid point about using AI to understand human language in the context of psychoactive substances. From a scientific standpoint, drug effects on human mental states have

been investigated through two primary approaches. The initial approach involves retrospective descriptive reports, which may be impacted by inaccurate recollections. These reports are typically non-standardized, making them more easily analyzed qualitatively, an analytic approach that can be time-consuming and lack generalizability. The application of AI for this type of analysis provides significant benefits. Another approach is automated speech-based technique to characterize clinically relevant alterations to mental state [203].

Computer-assisted cognitive rehabilitation systems and games for health are becoming increasingly common. To guarantee continuous, efficient, and long-term patient rehabilitation, these systems will need to be implemented within the home environment. However, current home systems often lack the presence of a therapist, which poses two major challenges. First, they require sensors and actuators that compensate for the absence of the therapist's eyes and hands. Second, these systems need to capture and utilize the therapist's expertise effectively [125].

On the other hand, robots equipped with machine learning capabilities have the advantage of being able to adapt their behavior and response mechanisms based on user input. This capability is particularly valuable when it comes to maximizing task performance and facilitating seamless social engagements. In the study [95], a robot directly asks a human user about their grasp of a learning objective, and adaptively adjusts the level of assistance it provides based on the response. Interestingly, the authors found that when the robot frequently queries the user for "learning-sensitive" information during every task-related question, users reported feeling more likely to rely on the robot's guidance.

In mental health, conversational AI has demonstrated both the potential to facilitate and inhibit the disclosure of information across various contexts. For instance, individuals have been found to be more open about reporting mental health symptoms when interacting with a conversational AI compared to a human listener, and these systems have successfully been implemented in treating persecutory delusions among individuals with psychosis [153, 204, 205].

Automated bots are currently unable to recreate the richness of a face-to-face interaction with a mental health professional, even with their efforts to mirror real-life interactions. Although some level of personalization is present (e.g., different tips or strategies are offered for individuals experiencing symptoms of depression vs. anxiety), the support provided remains generic and may be more akin to that of a self-help book at this stage [115].

A comparison of the effectiveness between artificial intelligence and human coaching in achieving goals was made in one of the other pieces of research. Research has demonstrated that AI can effectively replace human coaches who utilize simplistic, model-based coaching approaches. However, at the moment, AI lacks empathy and emotional intelligence, which are important aspects when it comes to effective coaching interventions [206]. Although most recent work claim that CVA outperforms humans in emotional awareness evaluations [207, 208].

One of the more perspective directions of development is the Explainable Artificial Intelligence (XAI). It represents a novel array of methodologies aimed at enhancing understanding and transparency in AI decision-making. In the context of neurostimulation, XAI holds significant value for both basic scientific research and therapeutic applications [71, 209–212].

In its current state, artificial intelligence does not possess the same level of intelligence as humans. Consequently, AI is unlikely to experience genuine mental illnesses as we understand them. Nevertheless, some experts posit that certain mental health concerns may emerge as computers become more advanced and interact more closely with humans. Potential problems AI may encounter encompass:

- Mental health issues: AI can function as a mirror to users, potentially becoming a source of stress, anxiety, and depression due to its interactions with them;
- Cognitive biases: AI may learn from and replicate human biases, including bigotry, prejudice, and discrimination [213].

One of the strengths of the reviewed experimental works in our study lies in their clear and comprehensive descriptions of the research procedures involved. Even studies with slightly flawed methodologies frequently offer detailed descriptions of their methodologies or cite previously published protocols as references for their methods. This clarity and transparency are crucial for understanding the research process and its inner workings, as well as for facilitating both replication and evaluation of the results.

A significant weakness in the field of research in the field of AI is the preponderance of pilot studies [117, 127, 200, 214–220]. While these initial studies play a crucial role in paving the way for future research, the majority of data generated from such studies is often not replicable or confirmable, leading to an overall decrease in research quality within the field. Small-scale studies are instrumental in the early stages of developing a technology or protocol, as they provide vital insights and data that can be utilized to refine the overall design. The absence of a control group in a study makes it challenging, if not impossible, to effectively control for any influence factors unrelated to the particular treatment being tested. This weakens the overall rigor and reliability of the study, making it difficult to accurately attribute any observed changes or effects solely to the intervention in question.

Another weakness is that AI system could be designed or trained poorly, or if it is given inaccurate or incomplete data, it may produce less accurate or reliable results. However, the development of AI systems requires a

combination of technical expertise, knowledge of algorithms and machine learning techniques, as well as access to large datasets and computing resources. Quality of markup on the train data should be improved for the better development in AI field.

It is essential to emphasize that our research has certain limitations. First, we acknowledge that we were unable to carry out a thorough assessment of each paper in relation to its respective field of interest, given the large sample size obtained from the PubMed database. Due to the sheer number of papers involved, it was impractical to manually review each one individually. While we made every effort to ensure thoroughness and accuracy in our review, some elements may have been overlooked. We attempted to address this challenge by applying several filters and inclusion criteria to generate relevant keywords for use in our search. While this method does not guarantee the complete exclusion of unsuitable papers from the final sample, there is a possibility that a few irrelevant articles may have been included. However, we can confidently state that the number of such papers is likely to be relatively small, and their inclusion did not significantly impact our overall results as a consequence of the rigorous screening process and the utilization of stringent filters and inclusion criteria.

5 Conclusions

Artificial intelligence has been effectively utilized for several years in aiding patients and healthcare professionals during rehabilitation processes. This article provides an examination of the current applications of AI methodologies within the field of mental rehabilitation. Functional connectivity analysis and virtual assistants are the most widely adopted form, yet new technologies are under investigation for their potential as diagnostic tools. Although the evidence supporting the utilization of AI in rehabilitation is encouraging, there is a paucity of systematic reviews that encompass a substantial body of work focusing on this topic. In addition, there is a necessity for extensive studies and systematic reviews examining various applications in diverse clinical populations.

This paper features a well-organized and straightforward way of presenting information, making it especially beneficial for specialists in the fields of AI, methods, and connected medical disciplines. It offers a clear overview of the various options available, detailing their advantages and drawbacks, thus serving as an informative guide or reference material for professionals seeking to utilize and apply distinct methods and techniques in their professional work.

Acknowledgements This work was supported by the Russian Ministry of Health as part of the scientific work “Development of a multimodal biofeedback-based hardware and software system for rehabilitation of patients with cognitive and motor disorders of different nature”, No. 123020600127-4, performed at the National Medical Research Center for Therapy and Preventive Medicine in 2023–2025.

Availability of data and materials The data presented in this study are available on request from the corresponding author.

References

1. Z. Bendella, C.N. Widmann, J.P. Layer, Y.L. Layer, R. Haase, M. Sauer, et al., Brain volume changes after Covid-19 compared to healthy controls by artificial intelligence-based MRI volumetry. *Diagnostics* (Basel, Switzerland) **13** (2023)
2. U. Hwang, S.W. Kim, D. Jung, S. Kim, H. Lee, S.W. Seo et al., Real-world prediction of preclinical Alzheimer’s disease with a deep generative model. *Artif. Intell. Med.* **144**, 102654 (2023)
3. M. Knop, S. Weber, M. Mueller, B. Niehaves, Human factors and technological characteristics influencing the interaction of medical professionals with artificial intelligence-enabled clinical decision support systems: literature review. *JMIR Hum. Factors* **9**, e28639 (2022)
4. C.Y. Tsai, Y.C. Kuan, W.H. Hsu, Y.T. Lin, C.R. Hsu, K. Lo, et al., Differentiation model for insomnia disorder and the respiratory arousal threshold phenotype in obstructive sleep apnea in the taiwanese population based on oximetry and anthropometric features. *Diagnostics* (Basel, Switzerland) **12** (2021)
5. C. Wachinger, M. Reuter, Domain adaptation for Alzheimer’s disease diagnostics. *Neuroimage* **139**, 470–479 (2016)
6. S.M. Andrade, L. da Silva-Sauer, C.D. de Carvalho, E.L.M. de Araújo, Ed.O. Lima, F.M.L. Fernandes et al., Identifying biomarkers for TDCS treatment response in Alzheimer’s disease patients: a machine learning approach using resting-state eeg classification. *Front. Hum. Neurosci.* **17**, 1234168 (2023)
7. L. Bajenaru, A. Sorici, I.G. Mocanu, A.M. Florea, F.A. Antochi, A.C. Ribigan, Shared decision-making to improve health-related outcomes for adults with stroke disease. *Healthcare* (Basel, Switzerland) **11** (2023)
8. A. Bishara, E.H. Maze, M. Maze, Considerations for the implementation of machine learning into acute care settings. *Br. Med. Bull.* **141**, 15–32 (2022)

9. L. Li, Y. Zhang, L. Huang, J. Zhao, J. Wang, T. Liu, Robot assisted treatment of hand functional rehabilitation based on visual motor imagination. *Front. Aging Neurosci.* **14**, 870871 (2022)
10. A. Mihailidis, M. Melonis, R. Keyfitz, M. Lanning, S. Van Vuuren, C. Bodine, A nonlinear contextually aware prompting system (n-caps) to assist workers with intellectual and developmental disabilities to perform factory assembly tasks: system overview and pilot testing. *Disabil. Rehabil. Assist. Technol.* **11**, 604–12 (2016)
11. A.R. Rahmanti, H.C. Yang, B.S. Bintoro, A.A. Nursetyo, M.S. Muhtar, S. Syed-Abdul et al., Slimme, a chatbot with artificial empathy for personal weight management: system design and finding. *Front. Nutr.* **9**, 870775 (2022). <https://doi.org/10.3389/fnut.2022.870775>
12. S. Rezaei, A. Moturu, S. Zhao, K.M. Prkachin, T. Hadjistavropoulos, B. Taati, Unobtrusive pain monitoring in older adults with dementia using pairwise and contrastive training. *IEEE J. Biomed. Health Inform.* **25**, 1450–1462 (2021)
13. R.S. Zoodsma, R. Bosch, T. Alderliesten, C.W. Bollen, T.H. Kappen, E. Koomen et al., Continuous data-driven monitoring in critical congenital heart disease: clinical deterioration model development. *JMIR Cardiol.* **7**, e45190 (2023)
14. C. Chandler, P.W. Foltz, B. Elvevåg, Improving the applicability of ai for psychiatric applications through human-in-the-loop methodologies. *Schizophr. Bull.* **48**(5), 949–957 (2022). <https://doi.org/10.1093/schbul/sbac038>
15. M. Ferrara, G. Franchini, M. Funaro, M. Cutroni, B. Valier, T. Toffanin et al., Machine learning and non-affective psychosis: identification, differential diagnosis, and treatment. *Curr. Psychiatry Rep.* **24**, 925–936 (2022)
16. S. Galderisi, G.M. Giordano, We are not ready to abandon the current schizophrenia construct, but should be prepared to do so. *Schizophr. Res.* **242**, 30–34 (2022)
17. W.S. Kim, D.W. Heo, J. Maeng, J. Shen, U. Tsogt, S. Odkhuu, et al., Deep learning-based brain age prediction in patients with schizophrenia spectrum disorders. *Schizophrenia Bull.* sbad167 (2023)
18. K.I. Okada, K. Miura, M. Fujimoto, K. Morita, M. Yoshida, H. Yamamori et al., Impaired inhibition of return during free-viewing behaviour in patients with schizophrenia. *Sci. Rep.* **11**, 3237 (2021)
19. G. Barbalat, J. Plasse, I. Chéreau-Boudet, B. Gouache, E. Legros-Lafarge, C. Massoubre et al., Contribution of socio-demographic and clinical characteristics to predict initial referrals to psychosocial interventions in patients with serious mental illness. *Epidemiol. Psychiatr. Sci.* **33**, e2 (2024)
20. A. Tkachev, E. Stekolshchikova, A. Vanyushkina, H. Zhang, A. Morozova, S. Zozulya et al., Lipid alteration signature in the blood plasma of individuals with schizophrenia, depression, and bipolar disorder. *JAMA Psychiatry* **80**, 250–259 (2023)
21. K.C. van den Berg, A.T. Hendrickson, S.A. Hales, M. Voncken, G.P.J. Keijsers, Comparing the effectiveness of imagery focussed cognitive therapy to group psychoeducation for patients with bipolar disorder: a randomised trial. *J. Affect. Disord.* **320**, 691–700 (2023)
22. Y.M. Chen, P.C. Chen, W.C. Lin, K.C. Hung, Y.C.B. Chen, C.F. Hung et al., Predicting new-onset post-stroke depression from real-world data using machine learning algorithm. *Front. Psychol.* **14**, 1195586 (2023)
23. E.N. Pitsik, V.A. Maximenko, S.A. Kurkin, A.P. Sergeev, D. Stoyanov, R. Paunova et al., The topology of fMRI-based networks defines the performance of a graph neural network for the classification of patients with major depressive disorder. *Chaos Solitons Fractals* **167**, 113041 (2023)
24. A.V. Andreev, S.A. Kurkin, D. Stoyanov, A.A. Badarin, R. Paunova, A.E. Hramov, Toward interpretability of machine learning methods for the classification of patients with major depressive disorder based on functional network measures. *Chaos Interdiscipl. J. Nonlinear Sci.* **33**(6) (2023)
25. C. Sinha, A.L. Cheng, M. Kadaba, Adherence and engagement with a cognitive behavioral therapy-based conversational agent (WYSA for chronic pain) among adults with chronic pain: survival analysis. *JMIR Form. Res.* **6**(5), e37302 (2022). <https://doi.org/10.2196/37302>
26. D. Stoyanov, V. Khorev, R. Paunova, S. Kandilarova, D. Simeonova, A. Badarin et al., Resting-state functional connectivity impairment in patients with major depressive episode. *Int. J. Environ. Res. Public Health* **19**(21), 14045 (2022)
27. H. Jin, S. Wu, P. Di Capua, Development of a clinical forecasting model to predict comorbid depression among diabetes patients and an application in depression screening policy making. *Prev. Chronic Dis.* **12**, E142 (2015)
28. M.K. van Vugt, P. Hitchcock, B. Shahar, W. Britton, The effects of mindfulness-based cognitive therapy on affective memory recall dynamics in depression: a mechanistic model of rumination. *Front. Hum. Neurosci.* **6**, 257 (2012)
29. M.S. Kabir, S. Kurkin, G. Portnova, O. Martynova, Z. Wang, A. Hramov, Contrastive machine learning reveals in EEG resting-state network salient features specific to autism spectrum disorder. *Chaos Solitons Fractals* **185**, 115123 (2024)
30. O. Sysoeva, V. Maximenko, A. Kuc, V. Voinova, O. Martynova, A. Hramov, Abnormal spectral and scale-free properties of resting-state EEG in girls with RETT syndrome. *Sci. Rep.* **13**(1), 12932 (2023)
31. N.A. Aleksandrova, A.E. Hramov, M.V. Khramova, Designing, implementation and use of robotic devices in the social sectors in foreign studies. In: 2018 IEEE International Conference “Quality Management, Transport and Information Security, Information Technologies” (IT &QM &IS) (IEEE, 2018) pp. 536–541
32. V.A. Maksimenko, A.E. Hramov, V.V. Grubov, V.O. Nedaivozov, V.V. Makarov, A.N. Pisarchik, Nonlinear effect of biological feedback on brain attentional state. *Nonlinear Dyn.* **95**(3), 1923–1939 (2019)
33. E.J. Topol, High-performance medicine: the convergence of human and artificial intelligence. *Nat. Med.* **25**(1), 44–56 (2019)

34. A.N. Pisarchik, V.A. Maksimenko, A.E. Hramov, From novel technology to novel applications: Comment on “an integrated brain–machine interface platform with thousands of channels” by Elon musk and Neuralink. *J. Med. Internet Res.* **21**(10), e16356 (2019)
35. A.E. Hramov, V.A. Maksimenko, A.N. Pisarchik, Physical principles of brain–computer interfaces and their applications for rehabilitation, robotics and control of human brain states. *Phys. Rep.* **918**, 1–133 (2021)
36. O.E. Karpov, V.V. Grubov, V.A. Maksimenko, S.A. Kurkin, N.M. Smirnov, N.P. Utyashev et al., Extreme value theory inspires explainable machine learning approach for seizure detection. *Sci. Rep.* **12**(1), 11474 (2022)
37. O.E. Karpov, M.S. Khoymov, V.A. Maksimenko, V.V. Grubov, N. Utyashev, D.A. Andrikov et al., Evaluation of unsupervised anomaly detection techniques in labelling epileptic seizures on human EEG. *Appl. Sci.* **13**(9), 5655 (2023)
38. M.A. Althobiani, Y. Ranjan, J. Jacob, M. Orini, R.J.B. Dobson, J.C. Porter et al., Evaluating a remote monitoring program for respiratory diseases: prospective observational study. *JMIR Form. Res.* **7**, e51507 (2023)
39. Z. Ahmed, K. Mohamed, S. Zeeshan, X. Dong, Artificial intelligence with multi-functional machine learning platform development for better healthcare and precision medicine. *Database* **2020**, baaa010 (2020)
40. M.M. Baig, H. GholamHosseini, A.A. Moqem, F. Mirza, M. Lindén, A systematic review of wearable patient monitoring systems—current challenges and opportunities for clinical adoption. *J. Med. Syst.* **41**, 115 (2017)
41. B.F. Kocyigit, M.I. Assylbek, M. Yessirkepov, Telerehabilitation: lessons from the Covid-19 pandemic and future perspectives. *Rheumatol. Int.* **44**(4), 577–582 (2024)
42. S. Graham, C. Depp, E.E. Lee, C. Nebeker, X. Tu, H.C. Kim et al., Artificial intelligence for mental health and mental illnesses: an overview. *Curr. Psychiatry Rep.* **21**, 1–18 (2019)
43. M. Tahan, T. Saleem, Application of artificial intelligence for diagnosis, prognosis and treatment in psychology: a review. *Neuropsychiatry i Neuropsychologia/Neuropsychiatry Neuropsychol.* **18**(1), 36–45 (2021)
44. J. Zhao, M. Wu, L. Zhou, X. Wang, J. Jia, Cognitive psychology-based artificial intelligence review. *Front. Neurosci.* **16**, 1024316 (2022)
45. A.E. Hramov, V.A. Maksimenko, S.V. Pchelintseva, A.E. Runnova, V.V. Grubov, V.Y. Musatov et al., Classifying the perceptual interpretations of a bistable image using EEG and artificial neural networks. *Front. Neurosci.* **11**, 674 (2017)
46. N. Kriegeskorte, P.K. Douglas, Cognitive computational neuroscience. *Nat. Neurosci.* **21**(9), 1148–1160 (2018)
47. S.M. Doane, Y.W. Sohn, Adapt: a predictive cognitive model of user visual attention and action planning. *User Model. User-Adap. Inter.* **10**, 1–45 (2000)
48. Y.T. Kim, H. Kim, C.H. Lee, B.C. Yoon, J.B. Kim, Y.H. Choi et al., Intracranial densitometry-augmented machine learning enhances the prognostic value of brain CT in pediatric patients with traumatic brain injury: a retrospective pilot study. *Front. Pediatr.* **9**, 750272 (2021)
49. S.M. LaConte, Decoding fMRI brain states in real-time. *Neuroimage* **56**, 440–54 (2011)
50. D.M. Lyall, A. Kormilitzin, C. Lancaster, J. Sousa, F. Petermann-Rocha, C. Buckley et al., Artificial intelligence for dementia-applied models and digital health. *Alzheimer’s Dementia* **19**, 5872–5884 (2023)
51. P. Mashouri, B. Taati, H. Quirt, A. Iaboni, Quality indicators as predictors of future inspection performance in Ontario nursing homes. *J. Am. Med. Dir. Assoc.* **21**, 793–798.e1 (2020)
52. V.A. Maksimenko, A.E. Runnova, M.O. Zhuravlev, P. Protasov, R. Kulanin, M.V. Khramova et al., Human personality reflects spatio-temporal and time-frequency EEG structure. *PLoS ONE* **13**(9), e0197642 (2018)
53. M. Maj, J. van Os, M. De Hert, W. Gaebel, S. Galderisi, M.F. Green et al., The clinical characterization of the patient with primary psychosis aimed at personalization of management. *World Psychiatry* **20**(1), 4–33 (2021)
54. H. Weerahandi, K.A. Hochman, E. Simon, C. Blaum, J. Chodosh, E. Duan et al., Post-discharge health status and symptoms in patients with severe Covid-19. *J. Gen. Intern. Med.* **36**, 738–745 (2021)
55. M.M. Adamson, S. Shakil, T. Sultana, M.A. Hasan, F. Mubarak, S.A. Enam et al., Brain injury and dementia in Pakistan: current perspectives. *Front. Neurol.* **11**, 299 (2020)
56. S.G. Ali, X. Wang, P. Li, Y. Jung, L. Bi, J. Kim et al., A systematic review: virtual-reality-based techniques for human exercises and health improvement. *Front. Public Health* (2023). <https://doi.org/10.3389/fpubh.2023.1143947>
57. M.F. Almufareh, S. Kausar, S. Humayunx, S. Tehsin, Leveraging motor imagery rehabilitation for individuals with disabilities: a review. *Healthcare (Basel, Switzerland)* **11**(19), 2653 (2023)
58. C.A. Goldstein, R.B. Berry, D.T. Kent, D.A. Kristo, A.A. Seixas, S. Redline et al., Artificial intelligence in sleep medicine: background and implications for clinicians. *J. Clin. Sleep Med.* **16**(4), 609–618 (2020). <https://doi.org/10.5664/jcsm.8388>
59. R. Bevilacqua, F. Barbarossa, L. Fantechi, D. Fornarelli, E. Paci, S. Bolognini, et al., Radiomics and artificial intelligence for the diagnosis and monitoring of Alzheimer’s disease: a systematic review of studies in the field. *J. Clin. Med.* **12**(16), 5432 (2023)
60. G. Brizzi, M. Sansoni, D. Di Lernia, F. Frisone, C. Tuena, G. Riva, The multisensory mind: a systematic review of multisensory integration processing in anorexia and bulimia nervosa. *J. Eat. Disord.* **11**, 204 (2023)
61. V. Catania, F. Rundo, S. Panerai, R. Ferri, Virtual reality for the rehabilitation of acquired cognitive disorders: a narrative review. *Bioengineering (Basel, Switzerland)* **11**(1), 35 (2023). <https://doi.org/10.3390/bioengineering11010035>
62. R. Charow, T. Jeyakumar, S. Younus, E. Dolatabadi, M. Salhia, D. Al-Mouaswas et al., Artificial intelligence education programs for health care professionals: scoping review. *JMIR Med. Educ.* **7**, e31043 (2021)

63. Y.J. Choo, M.C. Chang, Use of machine learning in stroke rehabilitation: a narrative review. *Brain Neurorehabil.* (2022). <https://doi.org/10.12786/bn.2022.15.e26>
64. Y. Cohen, T.A. Engel, C. Langdon, G.W. Lindsay, T. Ott, M.A.K. Peters et al., Recent advances at the interface of neuroscience and artificial neural networks. *J. Neurosci. Off. J. Soc. Neurosci.* **42**, 8514–8523 (2022)
65. J.L. Bender, S. Babinski, G. Wong, A.C. Tricco, M. Englesakis, A.B. Cyr et al., Establishing best practices in cancer online support groups: protocol for a realist review. *BMJ Open* **11**, e053916 (2021)
66. D.C. Billing, G.R. Fordy, K.E. Friedl, H. Hasselström, The implications of emerging technology on military human performance research priorities. *J. Sci. Med. Sport* **24**, 947–953 (2021)
67. C. Brown, R. Nazeer, A. Gibbs, P. Le Page, A.R. Mitchell, Breaking bias: the role of artificial intelligence in improving clinical decision-making. *Cureus* **15**(3) (2023)
68. L.J. Meier, A. Hein, K. Diepold, A. Buyx, Algorithms for ethical decision-making in the clinic: a proof of concept. *Am. J. Bioethics AJOB* **22**, 4–20 (2022)
69. J. Moreno, M.L. Gross, J. Becker, B. Hereth, N.D. Shortland, N.G. Evans, The ethics of AI-assisted warfighter enhancement research and experimentation: historical perspectives and ethical challenges. *Front. Big Data* **5**, 978734 (2022)
70. N.M. Safdar, J.D. Banja, C.C. Meltzer, Ethical considerations in artificial intelligence. *Eur. J. Radiol.* **122**, 108768 (2020)
71. J. van der Waa, S. Verdult, K. van den Bosch, J. van Diggelen, T. Haije, B. van der Stigchel et al., Moral decision making in human-agent teams: human control and the role of explanations. *Front. Robot. AI* **8**, 640647 (2021). <https://doi.org/10.3389/frobt.2021.640647>
72. L. Waltman, N.J. van Eck, E.C. Noyons, A unified approach to mapping and clustering of bibliometric networks. *J. Informetr.* **4**(4), 629–635 (2010). <https://doi.org/10.1016/j.joi.2010.07.002>
73. N.J. Van Eck, L. Waltman, Visualizing bibliometric networks, in *Measuring Scholarly Impact: Methods and Practice*. (Springer, Berlin, 2014), pp.285–320
74. O.E. Karpov, E.N. Pitsik, S.A. Kurkin, V.A. Maksimenko, A.V. Gusev, N.N. Shusharina et al., Analysis of publication activity and research trends in the field of ai medical applications: network approach. *Int. J. Environ. Res. Public Health* **20**(7), 5335 (2023)
75. L. Waltman, N.J. Van Eck, A smart local moving algorithm for large-scale modularity-based community detection. *Eur. Phys. J. B* **86**, 1–14 (2013)
76. F. Callaway, Y.R. Jain, B. van Opheusden, P. Das, G. Iwama, S. Gul et al., Leveraging artificial intelligence to improve people’s planning strategies. *Proc. Natl. Acad. Sci. USA* **119**(12), e2117432119 (2022). <https://doi.org/10.1073/pnas.2117432119>
77. S. Joksimovic, D. Ifenthaler, R. Marrone, M. De Laat, G. Siemens, Opportunities of artificial intelligence for supporting complex problem-solving: findings from a scoping review. *Comput. Educ. Artif. Intell.* **4**, 100138 (2023)
78. Y. Yao, J. Xu, J. Shi, B. Xu, Learning to activate logic rules for textual reasoning. *Neural Netw. Off. J. Int. Neural Netw. Soc.* **106**, 42–49 (2018)
79. M. Kopka, M.L. Schmieding, T. Rieger, E. Roesler, F. Balzer, M.A. Feufel, Determinants of laypersons’ trust in medical decision aids: Randomized controlled trial. *JMIR Hum. Factors* **9**, e35219 (2022)
80. T. Thomas, D. Straub, F. Tatai, M. Shene, T. Tosik, K. Kersting, et al., Modelling dataset bias in machine-learned theories of economic decision-making. *Nat. Hum. Behav.* **8**(4), 679–691 (2024)
81. L. Badino, A. D’Ausilio, L. Fadiga, G. Metta, Computational validation of the motor contribution to speech perception. *Top. Cogn. Sci.* **6**, 461–75 (2014)
82. Y. Wang, X. Zhang, C. Wang, W. Huang, Q. Xu, D. Liu et al., Modulation of biological motion perception in humans by gravity. *Nat. Commun.* **13**, 2765 (2022)
83. B. Grawemeyer, R. Cox, C. Lum, AUDIX: a knowledge-based system for speech-therapeutic auditory discrimination exercises. *Stud. Health Technol. Inform.* **77**, 568–72 (2000)
84. M. Krenn, R. Pollice, S.Y. Guo, M. Aldeghi, A. Cervera-Lierta, P. Friederich et al., On scientific understanding with artificial intelligence. *Nat. Rev. Phys.* **4**(12), 761–769 (2022). <https://doi.org/10.1038/s42254-022-00518-3>
85. M.J. Willemink, W.A. Koszek, C. Hardell, J. Wu, D. Fleischmann, H. Harvey et al., Preparing medical imaging data for machine learning. *Radiology* **295**, 4–15 (2020)
86. B. Tokar, M. Baskaya, O. Celik, F. Cemrek, A. Acikgoz, Application of machine learning techniques for enuresis prediction in children. *Eur. J. Pediatr. Surg. Off. J. Aust. Assoc. Pediatr. Surg Zeitschrift fur Kinderchirurgie* **31**, 414–419 (2021)
87. V.D. Veksler, B.E. Hoffman, N. Buchler, Symbolic deep networks: a psychologically inspired lightweight and efficient approach to deep learning. *Top. Cogn. Sci.* **14**, 702–717 (2022)
88. A.H. Thieme, Y. Zheng, G. Machiraju, C. Sadee, M. Mittermaier, M. Gertler et al., A deep-learning algorithm to classify skin lesions from MPOX virus infection. *Nat. Med.* **29**, 738–747 (2023)
89. B. Tahayori, N. Chini-Foroush, H. Akhlaghi, Advanced natural language processing technique to predict patient disposition based on emergency triage notes. *Emerg. Med. Australas. EMA* **33**, 480–484 (2021)
90. J.P. Funnell, K. Noor, D.Z. Khan, L. D’Antona, R.J.B. Dobson, J.G. Hanrahan, et al., Characterization of patients with idiopathic normal pressure hydrocephalus using natural language processing within an electronic healthcare record system. *J. Neurosurg.* **138**(6), 1731–1739 (2022)
91. Y. Gao, D. Dligach, T. Miller, J. Caskey, B. Sharma, M.M. Churpek et al., Dr. Bench: diagnostic reasoning benchmark for clinical natural language processing. *J. Biomed. Inform.* **138**, 104286 (2023)

92. S. Eppel, H. Xu, M. Bismuth, A. Aspuru-Guzik, Computer vision for recognition of materials and vessels in chemistry lab settings and the vector-labpics data set. *ACS Cent. Sci.* **6**, 1743–1752 (2020)
93. S. Mehdizadeh, E. Dolatabadi, K.D. Ng, A. Mansfield, A. Flint, B. Taati et al., Vision-based assessment of gait features associated with falls in people with dementia. *J. Gerontol. A Biol. Sci. Med. Sci.* **75**, 1148–1153 (2020)
94. J. Zhou, Q. Nguyen, S. Kamath, Y. Hacohen, C. Zhu, M.J. Fu et al., Hands to hexapods, wearable user interface design for specifying leg placement for legged robots. *Front. Robot. AI* **9**, 852270 (2022)
95. A. Langer, R. Feingold-Polak, O. Mueller, P. Kellmeyer, S. Levy-Tzedek, Trust in socially assistive robots: considerations for use in rehabilitation. *Neurosci. Biobehav. Rev.* **104**, 231–239 (2019). <https://doi.org/10.1016/j.neubiorev.2019.07.014>
96. A.M. Fazlollahi, M. Bakhtaidar, A. Alsayegh, R. Yilmaz, A. Winkler-Schwartz, N. Mirchi et al., Effect of artificial intelligence tutoring vs expert instruction on learning simulated surgical skills among medical students: a randomized clinical trial. *JAMA Netw. Open* **5**, e2149008 (2022)
97. D.D. Miller, E.W. Brown, Artificial intelligence in medical practice: the question to the answer? *Am. J. Med.* **131**, 129–133 (2018)
98. D.M. Abdullah, A.M. Abdulazeez, Machine learning applications based on SVM classification a review. *Qubahan Acad. J.* **1**(2), 81–90 (2021)
99. J.H. Lee, M. Marzelli, F.A. Jolesz, S.S. Yoo, Automated classification of fMRI data employing trial-based imagery tasks. *Med. Image Anal.* **13**, 392–404 (2009)
100. L. Nanni, M. Interlenghi, S. Brahnay, C. Salvatore, S. Papa, R. Nemni et al., Comparison of transfer learning and conventional machine learning applied to structural brain MRI for the early diagnosis and prognosis of Alzheimer's disease. *Front. Neurol.* **11**, 576194 (2020)
101. N. Dietz, V. Jaganathan, V. Alkin, J. Mettelle, M. Boakye, D. Drazin, Machine learning in clinical diagnosis, prognostication, and management of acute traumatic spinal cord injury (SCI): a systematic review. *J. Clin. Orthop. Trauma* **35**, 102046 (2022)
102. T. Zhou, K.H. Thung, M. Liu, F. Shi, C. Zhang, D. Shen, Multi-modal latent space inducing ensemble SVM classifier for early dementia diagnosis with neuroimaging data. *Med. Image Anal.* **60**, 101630 (2020)
103. A.E. Hramov, N.S. Frolov, V.A. Maksimenko, S.A. Kurkin, V.B. Kazantsev, A.N. Pisarchik, Functional networks of the brain: from connectivity restoration to dynamic integration. *Phys. Usp.* **64**(6), 584 (2021)
104. L. Zhang, Y. Li, L. Bian, Q. Luo, X. Zhang, B. Zhao, Cognitive impairment of patient with neurological cerebrovascular disease using the artificial intelligence technology guided by MRI. *Front. Public Health* **9**, 813641 (2021)
105. R. Savitha, S. Suresh, N. Sundararajan, A meta-cognitive learning algorithm for a fully complex-valued relaxation network. *Neural Netw. Off. J. Int. Neural Netw. Soc.* **32**, 209–18 (2012)
106. I.C. Chen, C.L. Chang, M.H. Chang, L.W. Ko, Atypical functional connectivity during rest and task-related dynamic alteration in young children with attention deficit hyperactivity disorder: an analysis using the phase-locking value. *Psychiatry Clin. Neurosci.* **76**, 235–245 (2022)
107. S. Zhang, H. Zhao, W. Wang, Z. Wang, X. Luo, A. Hramov et al., Edge-centric effective connection network based on multi-modal MRI for the diagnosis of Alzheimer's disease. *Neurocomputing* **552**, 126512 (2023)
108. N. Frolov, M.S. Kabir, V. Maksimenko, A. Hramov, Machine learning evaluates changes in functional connectivity under a prolonged cognitive load. *Chaos Interdiscipl. J. Nonlinear Sci.* **31**(10) (2021)
109. R. Islam, A.V. Andreev, N.N. Shusharina, A.E. Hramov, Explainable machine learning methods for classification of brain states during visual perception. *Mathematics* **10**(15), 2819 (2022)
110. C.M. Felmingham, N.R. Adler, Z. Ge, R.L. Morton, M. Janda, V.J. Mar, The importance of incorporating human factors in the design and implementation of artificial intelligence for skin cancer diagnosis in the real world. *Am. J. Clin. Dermatol.* **22**, 233–242 (2021)
111. K. Borys, Y.A. Schmitt, M. Nauta, C. Seifert, N. Krämer, C.M. Friedrich, et al., Explainable ai in medical imaging: an overview for clinical practitioners—beyond saliency-based XAI approaches. *Eur. J. Radiol.* **162**, 110786 (2023)
112. K.M. Hossain, M.A. Islam, S. Hossain, A. Nijholt, M.A.R. Ahad, Status of deep learning for EEG-based brain-computer interface applications. *Front. Comput. Neurosci.* **16**, 1006763 (2023)
113. A. Kuc, S. Korchagin, V.A. Maksimenko, N. Shusharina, A.E. Hramov, Combining statistical analysis and machine learning for EEG scalp topograms classification. *Front. Syst. Neurosci.* **15**, 716897 (2021)
114. A. Farzmahdi, K. Rajaei, M. Ghodrati, R. Ebrahimpour, S.M. Khaligh-Razavi, A specialized face-processing model inspired by the organization of monkey face patches explains several face-specific phenomena observed in humans. *Sci. Rep.* **6**, 25025 (2016)
115. K. Kretzschmar, H. Tyroll, G. Pavarini, A. Manzini, I. Singh, Can your phone be your therapist? Young people's ethical perspectives on the use of fully automated conversational agents (chatbots) in mental health support. *Biomed. Inform. Insights* **11**, 117822261982908 (2019). <https://doi.org/10.1177/1178222619829083>
116. J. Brinsley, B. Singh, C.A. Maher, A digital lifestyle program for psychological distress, wellbeing and return-to-work: a proof-of-concept study. *Arch. Phys. Med. Rehabil.* **104**, 1903–1912 (2023)
117. J.M. Noble, A. Zamani, M. Gharaat, D. Merrick, N. Maeda, A. Lambe Foster et al., Developing, implementing, and evaluating an artificial intelligence-guided mental health resource navigation chatbot for health care workers and their families during and following the Covid-19 pandemic: protocol for a cross-sectional study. *JMIR Res. Protoc.* **11**, e33717 (2022)

118. I.L. Yeh, J. Holst-Wolf, N. Elangovan, A.V. Cuppone, K. Lakshminarayan, L. Cappello et al., Effects of a robot-aided somatosensory training on proprioception and motor function in stroke survivors. *J. NeuroEng. Rehabil.* (2021). <https://doi.org/10.1186/s12984-021-00871-x>
119. S. Yang, R. Li, H. Li, K. Xu, Y. Shi, Q. Wang et al., Exploring the use of brain–computer interfaces in stroke neurorehabilitation. *Biomed. Res. Int.* **2021**, 9967348 (2021)
120. C.L. Chu, Y.P. Chen, C.C.P. Chen, C.K. Chen, H.N. Chang, C.H. Chang et al., Functional recovery patterns of hemorrhagic and ischemic stroke patients under post-acute care rehabilitation program. *Neuropsychiatr. Dis. Treat.* **16**, 1975–1985 (2020)
121. C. Zuo, J. Jin, E. Yin, R. Saab, Y. Miao, X. Wang et al., Novel hybrid brain–computer interface system based on motor imagery and p300. *Cogn. Neurodyn.* **14**, 253–265 (2020)
122. C. Urdiales, E.J. Pérez, G. Peinado, M. Fdez-Carmona, J.M. Peula, R. Annicchiarico et al., On the construction of a skill-based wheelchair navigation profile. *IEEE Trans. Neural Syst. Rehabil. Eng. Publ. IEEE Eng. Med. Biol. Soc.* **21**, 917–27 (2013)
123. R. Mane, Z. Wu, D. Wang, Poststroke motor, cognitive and speech rehabilitation with brain–computer interface: a perspective review. *Stroke Vasc. Neurol.* **7**(6), 541–549 (2022). <https://doi.org/10.1136/svn-2022-001506>
124. M. Schreuder, A. Riccio, M. Riseti, S. Dähne, A. Ramsay, J. Williamson et al., User-centered design in brain–computer interfaces—a case study. *Artif. Intell. Med.* **59**, 71–80 (2013)
125. M. Oliver, M.A. Teruel, J.P. Molina, D. Romero-Ayuso, P. González, Ambient intelligence environment for home cognitive telerehabilitation. *Sensors (Basel, Switzerland)* **18**(11), 3671 (2018). <https://doi.org/10.3390/s18113671>
126. D.W.K. Man, S.F. Tam, C.W.Y. Hui-Chan, Learning to live independently with expert systems in memory rehabilitation. *NeuroRehabilitation* **18**, 21–9 (2003)
127. P.W. Chen, N.A. Baune, I. Zwir, J. Wang, V. Swamidass, A.W.K. Wong, Measuring activities of daily living in stroke patients with motion machine learning algorithms: a pilot study. *Int. J. Environ. Res. Public Health* **18**(4), 1634 (2021)
128. Y.C. Kuan, L.K. Huang, Y.H. Wang, C.J. Hu, I.J. Tseng, H.C. Chen et al., Balance and gait performance in older adults with early-stage cognitive impairment. *Eur. J. Phys. Rehabil. Med.* **57**, 560–567 (2021)
129. M. Bocchetta, E.G. Todd, G. Peakman, D.M. Cash, R.S. Convery, L.L. Russell et al., Differential early subcortical involvement in genetic FTD within the GENFI cohort. *NeuroImage Clin.* **30**, 102646 (2021)
130. S.Y. Yang, H.C. Lee, C.M. Huang, J.J. Chen, Efficacy of Tai Chi-style multi-component exercise on frontal-related cognition and physical health in elderly with amnesic mild cognitive impairment. *Front. Aging* **2**, 636390 (2021)
131. E. Rauseo, A. Salih, Z. Raisi-Estabragh, N. Aung, N. Khanderia, G.G. Slabaugh et al., Ischemic heart disease and vascular risk factors are associated with accelerated brain aging. *JACC Cardiovasc. Imaging* **16**, 905–915 (2023)
132. J.M. Pavon, L. Preville, M. Woo, R. Henao, M. Solomon, U. Rogers et al., Machine learning functional impairment classification with electronic health record data. *J. Am. Geriatr. Soc.* **71**, 2822–2833 (2023)
133. S. Sarraf, D.D. Desouza, J. Anderson, C. Saverino, MCADNNet: recognizing stages of cognitive impairment through efficient convolutional FMRI and MRI neural network topology models. *IEEE Access Pract. Innov. Open Solut.* **7**, 155584–155600 (2019)
134. E. Dolatabadi, Y.X. Zhi, A.J. Flint, A. Mansfield, A. Iaboni, B. Taati, The feasibility of a vision-based sensor for longitudinal monitoring of mobility in older adults with dementia. *Arch. Gerontol. Geriatr.* **82**, 200–206 (2019)
135. C. Tuena, C. Pupillo, C. Stramba-Badiale, M. Stramba-Badiale, G. Riva, Predictive power of gait and gait-related cognitive measures in amnesic mild cognitive impairment: a machine learning analysis. *Front. Hum. Neurosci.* **17**, 1328713 (2023)
136. A. Akl, B. Taati, A. Mihailidis, Autonomous unobtrusive detection of mild cognitive impairment in older adults. *IEEE Trans. Biomed. Eng.* **62**, 1383–1394 (2015)
137. D. Zhou, Y. Jin, Y. Chen, The application scenarios study on the intervention of cognitive decline in elderly population using metaverse technology. *Sheng wu yi xue gong cheng xue za zhi J. Biomed. Eng. Shengwu yixue gongchengxue zazhi* **40**, 573–581 (2023)
138. M. Malcangi, AI-based methods and technologies to develop wearable devices for prosthetics and predictions of degenerative diseases. *Methods Mol. Biol. (Clifton, NJ)* **2190**, 337–354 (2021)
139. B. Ma, J. Yang, F.K.Y. Wong, A.K.C. Wong, T. Ma, J. Meng et al., Artificial intelligence in elderly healthcare: a scoping review. *Ageing Res. Rev.* **83**, 101808 (2023)
140. S. Mehdizadeh, K. Van Ooteghem, H. Gulka, H. Nabavi, M. Faieghi, B. Taati et al., A systematic review of center of pressure measures to quantify gait changes in older adults. *Exp. Gerontol.* **143**, 111170 (2021)
141. A.L. Young, R.V. Marinescu, N.P. Oxtoby, M. Bocchetta, K. Yong, N.C. Firth et al., Uncovering the heterogeneity and temporal complexity of neurodegenerative diseases with subtype and stage inference. *Nat. Commun.* (2018). <https://doi.org/10.1038/s41467-018-05892-0>
142. A. Billot, S. Lai, M. Varkanitsa, E.J. Braun, B. Rapp, T.B. Parrish et al., Multimodal neural and behavioral data predict response to rehabilitation in chronic poststroke aphasia. *Stroke* **53**(5), 1606–1614 (2022). <https://doi.org/10.1161/strokeaha.121.036749>
143. A. García-Rudolph, A. García-Molina, E. Opisso, J.M. Tormos, V.I. Madai, D. Frey et al., Neuropsychological assessments of patients with acquired brain injury: a cluster analysis approach to address heterogeneity in web-based cognitive rehabilitation. *Front. Neurol.* (2021). <https://doi.org/10.3389/fneur.2021.701946>
144. L. Laranjo, A.G. Dunn, H.L. Tong, A.B. Kocaballi, J. Chen, R. Bashir et al., Conversational agents in healthcare: a systematic review. *J. Am. Med. Inform. Assoc.* **25**(9), 1248–1258 (2018). <https://doi.org/10.1093/jamia/ocy072>

145. L.C. MacLagan, M. Abdalla, D.A. Harris, T.A. Stukel, B. Chen, E. Candido et al., Can patients with dementia be identified in primary care electronic medical records using natural language processing? *J. Healthc. Inform. Res.* **7**, 42–58 (2023)
146. V.C. Kaelin, M. Valizadeh, Z. Salgado, J.G. Sim, D. Anaby, A.D. Boyd et al., Capturing and operationalizing participation in pediatric re/habilitation research using artificial intelligence: a scoping review. *Front. Rehabil. Sci.* (2022). <https://doi.org/10.3389/fresc.2022.855240>
147. S. Jirayucharoensak, P. Israsena, S. Pan-Ngum, S. Hemrungrojn, M. Maes, A game-based neurofeedback training system to enhance cognitive performance in healthy elderly subjects and in patients with amnesic mild cognitive impairment. *Clin. Interv. Aging* **14**, 347–360 (2019)
148. L. Borriore, H. Bellini, L.B. Razza, A.G. Avila, C. Baeken, A.K. Brem et al., Precision non-implantable neuromodulation therapies: a perspective for the depressed brain. *Braz. J. Psychiatry* **42**(4), 403–419 (2020). <https://doi.org/10.1590/1516-4446-2019-0741>
149. C. Vidaurre, S. Haufe, T. Jorajuría, K.R. Müller, V.V. Nikulin, Sensorimotor functional connectivity: a neurophysiological factor related to BCI performance. *Front. Neurosci.* **14**, 575081 (2020)
150. J. Chen, S. Zhao, H. Meng, X. Cheng, W. Tan, An interactive game for rehabilitation based on real-time hand gesture recognition. *Front. Physiol.* **13**, 1028907 (2022)
151. D. Mizrahi, I. Zuckerman, I. Laufer, Using a stochastic agent model to optimize performance in divergent interest tacit coordination games. *Sensors (Basel, Switzerland)* **20**(24), 7026 (2020). <https://doi.org/10.3390/s20247026>
152. S. Greer, D. Ramo, Y.J. Chang, M. Fu, J. Moskowicz, J. Haritatos, Use of the chatbot “vivibot” to deliver positive psychology skills and promote well-being among young people after cancer treatment: randomized controlled feasibility trial. *JMIR Mhealth Uhealth* **7**(10), e15018 (2019). <https://doi.org/10.2196/15018>
153. M. Danieli, T. Ciulli, S.M. Mousavi, G. Riccardi, A conversational artificial intelligence agent for a mental health care app: evaluation study of its participatory design. *JMIR Form. Res.* **5**(12), e30053 (2021). <https://doi.org/10.2196/30053>
154. T.A. Creed, L. Salama, R. Slevin, M. Tanana, Z. Imel, S. Narayanan et al., Enhancing the quality of cognitive behavioral therapy in community mental health through artificial intelligence generated fidelity feedback (project affect): a study protocol. *BMC Health Serv. Res.* **22**(1), 1177 (2022). <https://doi.org/10.1186/s12913-022-08519-9>
155. Y. He, L. Yang, C. Qian, T. Li, Z. Su, Q. Zhang et al., Conversational agent interventions for mental health problems: systematic review and meta-analysis of randomized controlled trials. *J. Med. Internet Res.* **25**, e43862 (2023)
156. D.E. Fleck, N. Ernest, C.M. Adler, K. Cohen, J.C. Eliassen, M. Norris et al., Prediction of lithium response in first-episode mania using the lithium intelligent agent (LITHIA): pilot data and proof-of-concept. *Bipolar Disord.* **19**(4), 259–272 (2017). <https://doi.org/10.1111/bdi.12507>
157. B. Inkster, S. Sarda, V. Subramanian, An empathy-driven, conversational artificial intelligence agent (WYSA) for digital mental well-being: real-world data evaluation mixed-methods study. *JMIR Mhealth Uhealth* **6**(11), e12106 (2018). <https://doi.org/10.2196/12106>
158. S. Raffard, R.N. Salesse, C. Bortolon, B.G. Bardy, J. Henriques, L. Marin et al., Using mimicry of body movements by a virtual agent to increase synchronization behavior and rapport in individuals with schizophrenia. *Sci. Rep.* (2018). <https://doi.org/10.1038/s41598-018-35813-6>
159. R. Liu, J.P. Salisbury, A. Vahabzadeh, N.T. Sahin, Feasibility of an autism-focused augmented reality smartglasses system for social communication and behavioral coaching. *Front. Pediatr.* (2017). <https://doi.org/10.3389/fped.2017.00145>
160. A. Vahabzadeh, N. Keshav, R. Abdus-Sabur, K. Huey, R. Liu, N. Sahin, Improved socio-emotional and behavioral functioning in students with autism following school-based smartglasses intervention: multi-stage feasibility and controlled efficacy study. *Behav. Sci.* **8**(10), 85 (2018). <https://doi.org/10.3390/bs8100085>
161. Q. Zheng, F. Liu, S. Xu, J. Hu, H. Lu, T. Liu, Artificial intelligence empowering research on loneliness, depression and anxiety-using Covid-19 as an opportunity. *J. Saf. Sci. Resil.* (2023)
162. B. Pani, J. Crawford, K.A. Allen, Can generative artificial intelligence foster belongingness, social support, and reduce loneliness? A conceptual analysis. *Appl. Gener. AI* 261–276 (2024)
163. K. Loveys, G. Fricchione, K. Kolappa, M. Sagar, E. Broadbent, Reducing patient loneliness with artificial agents: design insights from evolutionary neuropsychiatry. *J. Med. Internet Res.* **21**(7), e13664 (2019). <https://doi.org/10.2196/13664>
164. S. Han, C. Park, J.S.H. You, Effects of robotic interactive gait training combined with virtual reality and augmented reality on balance, gross motor function, gait kinetic, and kinematic characteristics in Angelman syndrome: a case report. *Children* **9**(4), 544 (2022). <https://doi.org/10.3390/children9040544>
165. H. Chang, S.H. Woo, S. Kang, C.Y. Lee, J.Y. Lee, J.K. Ryu, A curtailed task for quantitative evaluation of visuomotor adaptation in the head-mounted display virtual reality environment. *Front. Psychol.* **13**, 963303 (2022). <https://doi.org/10.3389/fpsy.2022.963303>
166. A. Tacchino, J. Podda, V. Bergamaschi, L. Pedullà, G. Bricchetto, Cognitive rehabilitation in multiple sclerosis: three digital ingredients to address current and future priorities. *Front. Hum. Neurosci.* **17**, 1130231 (2023). <https://doi.org/10.3389/fnhum.2023.1130231>
167. I.M. Chen, Y.Y. Chen, S.C. Liao, Y.H. Lin, Development of digital biomarkers of mental illness via mobile apps for personalized treatment and diagnosis. *J. Personal. Med.* **12**(6), 936 (2022)

168. S. Berrouiguet, M.L. Barrigón, J.L. Castroman, P. Courtet, A. Artés-Rodríguez, E. Baca-García, Combining mobile-health (Mhealth) and artificial intelligence (AI) methods to avoid suicide attempts: the smartcrises study protocol. *BMC Psychiatry* (2019). <https://doi.org/10.1186/s12888-019-2260-y>
169. C. Voss, J. Schwartz, J. Daniels, A. Kline, N. Haber, P. Washington et al., Effect of wearable digital intervention for improving socialization in children with autism spectrum disorder: a randomized clinical trial. *JAMA Pediatr.* **173**(5), 446 (2019). <https://doi.org/10.1001/jamapediatrics.2019.0285>
170. R. Medina, J. Bouhaben, I. de Ramón, P. Cuesta, L. Antón-Toro, J. Pacios et al., Electrophysiological brain changes associated with cognitive improvement in a pediatric attention deficit hyperactivity disorder digital artificial intelligence-driven intervention: randomized controlled trial. *J. Med. Internet Res.* **23**(11), e25466 (2021). <https://doi.org/10.2196/25466>
171. J. Tani, Y.H. Yang, C.M. Chen, C.Y. Siow, T.S. Chang, K. Yang et al., Domain-specific cognitive prosthesis for face memory and recognition. *Diagnostics* **12**(9), 2242 (2022). <https://doi.org/10.3390/diagnostics12092242>
172. K. White, S. Tate, R. Zafonte, S. Narayanan, M.R. Mehl, M. Shin et al., Socialbit: protocol for a prospective observational study to validate a wearable social sensor for stroke survivors with diverse neurological abilities. *BMJ Open* **13**(8), e076297 (2023). <https://doi.org/10.1136/bmjopen-2023-076297>
173. V. Vakili, W.P. Brinkman, M.A. Neerincx, Lessons learned from the development of technological support for PTSD prevention: a review. *Stud. Health Technol. Inform.* **181**, 22–6 (2012)
174. E.N. Pitsik, N.S. Frolov, N. Shusharina, A.E. Hramov, Age-related changes in functional connectivity during the sensorimotor integration detected by artificial neural network. *Sensors* **22**(7), 2537 (2022)
175. N. Frolov, V. Maksimenko, A. Lüttjohann, A. Koronovskii, A. Hramov, Feed-forward artificial neural network provides data-driven inference of functional connectivity. *Chaos Interdiscipl. J. Nonlinear Sci.* **29**(9) (2019)
176. A.E. Hramov, N.S. Frolov, V.A. Maksimenko, V.V. Makarov, A.A. Koronovskii, J. Garcia-Prieto, et al., Artificial neural network detects human uncertainty. *Chaos Interdiscipl. J. Nonlinear Sci.* **28**(3) (2018)
177. F. Yuan, E. Klavon, Z. Liu, R.P. Lopez, X. Zhao, A systematic review of robotic rehabilitation for cognitive training. *Front. Robot. AI* (2021). <https://doi.org/10.3389/frobt.2021.605715>
178. Y. Yang, The talent training mode of international service design using a human–computer interaction intelligent service robot from the perspective of cognitive psychology. *Front. Psychol.* **12**, 600218 (2021). <https://doi.org/10.3389/fpsyg.2021.600218>
179. Y. Wang, H. Zhu, N. Elangovan, L. Cappello, G. Sandini, L. Masia et al., A robot-aided visuomotor wrist training induces gains in proprioceptive and movement accuracy in the contralateral wrist. *Sci. Rep.* (2021). <https://doi.org/10.1038/s41598-021-84767-9>
180. B. Sawik, S. Tobis, E. Baum, A. Suwalska, S. Kropińska, K. Stachnik et al., Robots for elderly care: review, multi-criteria optimization model and qualitative case study. *Healthcare (Basel, Switzerland)* **11**(9), 1286 (2023). <https://doi.org/10.3390/healthcare11091286>
181. M. Simonov, G. Delconte, Humanoid assessing rehabilitative exercises. *Methods Inf. Med.* **54**, 114–21 (2015)
182. V. Sharma, M. Gupta, A. Kumar, D. Mishra, Video processing using deep learning techniques: a systematic literature review. *IEEE Access* **9**, 139489–139507 (2021)
183. S.Y. Gordleeva, S.A. Lobov, N.A. Grigorev, A.O. Savosenkov, M.O. Shamshin, M.V. Lukoyanov et al., Real-time EEG-EMG human–machine interface-based control system for a lower-limb exoskeleton. *IEEE Access* **8**, 84070–84081 (2020)
184. P. Chholak, G. Niso, V.A. Maksimenko, S.A. Kurkin, N.S. Frolov, E.N. Pitsik et al., Visual and kinesthetic modes affect motor imagery classification in untrained subjects. *Sci. Rep.* **9**(1), 9838 (2019)
185. H.L. Halme, L. Parkkonen, Comparing features for classification of MEG responses to motor imagery. *PLoS ONE* **11**(12), e0168766 (2016)
186. H. Altaheri, G. Muhammad, M. Alsulaiman, S.U. Amin, G.A. Altuwaijri, W. Abdul et al., Deep learning techniques for classification of electroencephalogram (EEG) motor imagery (MI) signals: a review. *Neural Comput. Appl.* **35**(20), 14681–14722 (2023)
187. E. Pitsik, N. Frolov, K. Hauke Kraemer, V. Grubov, V. Maksimenko, J. Kurths, et al., Motor execution reduces EEG signals complexity: Recurrence quantification analysis study. *Chaos Interdiscipl. J. Nonlinear Sci.* **30**(2) (2020)
188. V.A. Maksimenko, S.A. Kurkin, E.N. Pitsik, V.Y. Musatov, A.E. Runnova, T.Y. Efremova et al., Artificial neural network classification of motor-related EEG: an increase in classification accuracy by reducing signal complexity. *Complexity* **2018**(1), 9385947 (2018)
189. S. Kurkin, A. Badarin, V. Grubov, V. Maksimenko, A. Hramov, The oxygen saturation in the primary motor cortex during a single hand movement: functional near-infrared spectroscopy (FNIRS) study. *Eur. Phys. J. Plus* **136**(5), 548 (2021)
190. H. Chhabra, N. Shajil, G. Venkatasubramanian et al., Investigation of deep convolutional neural network for classification of motor imagery FNIRS signals for BCI applications. *Biomed. Signal Process. Control* **62**, 102133 (2020)
191. A.E. Hramov, V. Grubov, A. Badarin, V.A. Maksimenko, A.N. Pisarchik, Functional near-infrared spectroscopy for the classification of motor-related brain activity on the sensor-level. *Sensors* **20**(8), 2362 (2020)
192. H. Wang, Q. Su, Z. Yan, F. Lu, Q. Zhao, Z. Liu et al., Rehabilitation treatment of motor dysfunction patients based on deep learning brain–computer interface technology. *Front. Neurosci.* **14**, 595084 (2020)
193. A. Chepurova, A. Hramov, S. Kurkin, Motor imagery: how to assess, improve its performance, and apply it for psychosis diagnostics. *Diagnostics* **12**(4), 949 (2022)

194. N.S. Frolov, E.N. Pitsik, V.A. Maksimenko, V.V. Grubov, A.R. Kiselev, Z. Wang et al., Age-related slowing down in the motor initiation in elderly adults. *PLoS ONE* **15**(9), e0233942 (2020)
195. A. Cristiano, S. Musteata, S. De Silvestri, V. Bellandi, P. Ceravolo, M. Cesari et al., Older adults' and clinicians' perspectives on a smart health platform for the aging population: design and evaluation study. *JMIR Aging* **5**, e29623 (2022)
196. A. Garcia-Rudolph, E. Opisso, J.M. Tormos, V.I. Madai, D. Frey, H. Becerra et al., Toward personalized web-based cognitive rehabilitation for patients with ischemic stroke: Elo rating approach. *JMIR Med. Inform.* **9**(11), e28090 (2021). <https://doi.org/10.2196/28090>
197. A. Singh, S. Chakraborty, Z. He, S. Tian, S. Zhang, M.L.A. Lustria et al., Deep learning-based predictions of older adults' adherence to cognitive training to support training efficacy. *Front. Psychol.* **13**, 980778 (2022). <https://doi.org/10.3389/fpsyg.2022.980778>
198. S.B. Tan, J. Tan, M.N. Raczkowska, J. Chean Wen Lee, B. Rai, A. Remus et al., Digital game-based interventions for cognitive training in healthy adults and adults with cognitive impairment: protocol for a two-part systematic review and meta-analysis. *BMJ Open* **13**(5), e071059 (2023). <https://doi.org/10.1136/bmjopen-2022-071059>
199. P. Urwyler, R.K. Gupta, M. Falkner, J. Niklaus, R.M. Müri, T. Nef, Tablet-based puzzle game intervention for cognitive function and well-being in healthy adults: pilot feasibility randomized controlled trial. *JMIR Aging* **6**, e46177–e46177 (2023). <https://doi.org/10.2196/46177>
200. S.J. Eun, E.J. Kim, J.Y. Kim, Development and evaluation of an artificial intelligence-based cognitive exercise game: a pilot study. *J. Environ. Public Health* **2022**, 1–15 (2022). <https://doi.org/10.1155/2022/4403976>
201. D. Tolks, J.J. Schmidt, S. Kuhn, The role of AI in serious games and gamification for health: scoping review. *JMIR Serious Games* **12**, e48258 (2024). <https://doi.org/10.2196/48258>
202. G. Bedi, G.A. Cecchi, D.F. Slezak, F. Carrillo, M. Sigman, H. De Wit, A window into the intoxicated mind? Speech as an index of psychoactive drug effects. *Neuropsychopharmacology* **39**(10), 2340–2348 (2014)
203. P. Gaikwad, M. Venkatesan, Speech recognition-based prediction for mental health and depression: a review. In: *Congress on Control, Robotics, and Mechatronics* (Springer, 2023), pp. 13–24
204. A.S. Miner, N. Shah, K.D. Bullock, B.A. Arnow, J. Bailenson, J. Hancock, Key considerations for incorporating conversational ai in psychotherapy. *Front. Psychiatry* (2019). <https://doi.org/10.3389/fpsy.2019.00746>
205. T.K. Craig, M. Rus-Calafell, T. Ward, J.P. Leff, M. Huckvale, E. Howarth et al., Avatar therapy for auditory verbal hallucinations in people with psychosis: a single-blind, randomised controlled trial. *Lancet Psychiatry* **5**(1), 31–40 (2018)
206. N. Terblanche, J. Molyneux, E. de Haan, V.O. Nilsson, Comparing artificial intelligence and human coaching goal attainment efficacy. *PLoS ONE* **17**(6), e0270255 (2022). <https://doi.org/10.1371/journal.pone.0270255>
207. Z. Elyoseph, D. Hadar-Shoval, K. Asraf, M. Lvovsky, Chatgpt outperforms humans in emotional awareness evaluations. *Front. Psychol.* **14**, 1199058 (2023)
208. A.L. Faria, Y. Almeida, D. Branco, J. Câmara, M. Cameirão, L. Ferreira et al., Neuroaireh@b: an artificial intelligence-based methodology for personalized and adaptive neurorehabilitation. *Front. Neurol.* **14**, 1258323 (2023). <https://doi.org/10.3389/fneur.2023.1258323>
209. J.M. Fellous, G. Sapiro, A. Rossi, H. Mayberg, M. Ferrante, Explainable artificial intelligence for neuroscience: behavioral neurostimulation. *Front. Neurosci.* (2019). <https://doi.org/10.3389/fnins.2019.01346>
210. I. Chraïbi Kaadoud, A. Bennetot, B. Mawhin, V. Charisi, N. Díaz-Rodríguez, Explaining aha! moments in artificial agents through IKE-XAI: implicit knowledge extraction for explainable AI. *Neural Netw. Off. J. Int. Neural Netw. Soc.* **155**, 95–118 (2022)
211. S. Coppolino, M. Migliore, An explainable artificial intelligence approach to spatial navigation based on hippocampal circuitry. *Neural Netw. Off. J. Int. Neural Netw. Soc.* **163**, 97–107 (2023)
212. O.E. Karpov, S. Afinogenov, V.V. Grubov, V. Maksimenko, S. Korchagin, N. Utyashev et al., Detecting epileptic seizures using machine learning and interpretable features of human EEG. *Eur. Phys. J. Spec. Top.* **232**(5), 673–682 (2023)
213. H. Ashrafiyan, Can artificial intelligences suffer from mental illness? A philosophical matter to consider. *Sci. Eng. Ethics* **23**(2), 403–412 (2016). <https://doi.org/10.1007/s11948-016-9783-0>
214. H. Nabavi, S. Mehdizadeh, L.C. Shum, A.J. Flint, A. Mansfield, B. Taati et al., A pilot observational study of gait changes over time before and after an unplanned hospital visit in long-term care residents with dementia. *BMC Geriatr.* **23**, 723 (2023)
215. B.W. Heller, D. Datta, J. Howitt, A pilot study comparing the cognitive demand of walking for transfemoral amputees using the intelligent prosthesis with that using conventionally damped knees. *Clin. Rehabil.* **14**, 518–22 (2000)
216. C.Y. Tsai, S.M. Wu, Y.C. Kuan, Y.T. Lin, C.R. Hsu, W.H. Hsu et al., Associations between risk of Alzheimer's disease and obstructive sleep apnea, intermittent hypoxia, and arousal responses: a pilot study. *Front. Neurol.* **13**, 1038735 (2022)
217. Z. Zheng, Z. Warren, A. Weitlauf, Q. Fu, H. Zhao, A. Swanson et al., Brief report: evaluation of an intelligent learning environment for young children with autism spectrum disorder. *J. Autism Dev. Disord.* **46**, 3615–3621 (2016)
218. W.T. Liu, H.T. Huang, H.Y. Hung, S.Y. Lin, W.H. Hsu, F.Y. Lee, et al., Continuous positive airway pressure reduces plasma neurochemical levels in patients with OSA: a pilot study. *Life (Basel, Switzerland)* **13**(3), 613 (2023)

219. N. Hird, T. Osaki, S. Ghosh, S.K. Palaniappan, K. Maeda, Enabling personalization for digital cognitive stimulation to support communication with people with dementia: pilot intervention study as a prelude to AI development. *JMIR Form. Res.* **8**, e51732 (2024)
220. B.L. Edlow, Y.G. Bodien, T. Baxter, H.G. Belanger, R.J. Cali, K.B. Deary et al., Long-term effects of repeated blast exposure in united states special operations forces personnel: a pilot study protocol. *J. Neurotrauma* **39**, 1391–1407 (2022)

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.